

MIXING ACTIONS OF THE RATIONALS

RICHARD MILES AND TOM WARD

ABSTRACT. We study mixing properties of algebraic actions of $\mathbb{Q}^d$, showing in particular that prime mixing $\mathbb{Q}^d$ actions on connected groups are mixing of all orders, as is the case for $\mathbb{Z}^d$ -actions. This is shown using a uniform result on the solution of S -unit equations in characteristic zero fields due to Evertse, Schlickewei and Schmidt. In contrast, algebraic actions of the much larger group $\mathbb{Q}^*$ are shown to behave quite differently, with finite order of mixing possible on connected groups.

Mixing properties of $\mathbb{Z}^d$ -actions by automorphisms of a compact metrizable abelian group are quite well understood. Roughly speaking, the picture has three facets. Firstly, the correspondence between such actions and countably generated modules over the integral group ring $R_d = \mathbb{Z}[\mathbb{Z}^d]$ of the acting group $\mathbb{Z}^d$ due to Kitchens and Schmidt [6] allows any mixing problem to be reduced to the case corresponding to a cyclic module of the form R_d/P for a prime ideal $P \subset R_d$. Secondly, in the connected case $P \cap \mathbb{Z} = \{0\}$, Schmidt and Ward [12] showed that mixing implies mixing of all orders by relating the mixing property to S -unit equations and exploiting a deep result of Schlickewei on solutions of such equations [10] (see also [3] and [13]). Finally, in the totally disconnected case $P \cap \mathbb{Z} = p\mathbb{Z}$ for some rational prime p , Masser [9] has shown that the order of mixing is determined by the mixing behaviour of shapes, reducing the problem – in principle – to an algebraic one.

Our purpose here is to show how some of this changes for algebraic actions of infinitely generated abelian groups. The algebra is more involved, so for simplicity we restrict attention to the simplest extreme examples: actions of $\mathbb{Q}_{>0}^\times$ (isomorphic to the direct sum of countably many copies of $\mathbb{Z}$) and actions of $\mathbb{Q}^d$ (which is a torsion extension of $\mathbb{Z}^d$). These groups are the simplest non-trivial examples chosen from the ‘dual’ categories of free abelian and infinitely divisible groups in the sense of MacLane [8]. The algebraic difficulties mean we cannot present the complete picture found for $\mathbb{Z}^d$ -actions, and the emphasis is partly on revealing or suggestive examples.

Let α be an action of a countable abelian group Γ on a probability space $(X, \mathcal{B}, \mu)$. For a sequence (γ_n) in Γ , write $\gamma_n \rightarrow \infty$ if for every finite set $F \subset \Gamma$ there is an N for which $n > N$ implies that $\gamma_n \notin F$.

The action α is said to be *mixing on r sets* if for any sets

$$A_1, \dots, A_r \in \mathcal{B},$$

$$\mu(\alpha_{\gamma_1} A_1 \cap \dots \cap \alpha_{\gamma_r} A_r) \rightarrow \mu(A_1) \cdots \mu(A_r) \text{ as } \gamma_s - \gamma_t \rightarrow \infty \text{ for } s \neq t.$$

The *order of mixing* $\mathcal{M}(\alpha)$ of α is the largest value of r for which α is mixing on r sets, and α is said to be *mixing of all orders*, denoted $\mathcal{M}(\alpha) = \infty$, if it is mixing on r sets for all r .

1. ALGEBRAIC ACTIONS

Just as for algebraic $\mathbb{Z}^d$ -actions (see Schmidt's monograph [11]), Pontryagin duality gives a description of Γ -actions by automorphisms of compact abelian groups in terms of modules over the ring $\mathbb{Z}[\Gamma]$. If α is a Γ -action by automorphisms of X , then the character group $M = \widehat{X}$ inherits the structure of a $\mathbb{Z}[\Gamma]$ -module via the dual automorphisms $\widehat{\alpha}_\gamma$ for $\gamma \in \Gamma$; conversely any $\mathbb{Z}[\Gamma]$ -module M defines a compact abelian group $X_M = \widehat{M}$ carrying a dual Γ -action α_M . Write $\lambda = \lambda_X$ for the Haar measure on X .

A module is called *cyclic* if it is singly generated as a module, so takes the form $\mathbb{Z}[\Gamma]/I$ for some ideal $I \subset \mathbb{Z}[\Gamma]$, and the dual Γ -action will be called *prime* (or *radical*) if the module takes the form $\mathbb{Z}[\Gamma]/P$ for some prime (resp. radical) ideal $P \subset \mathbb{Z}[\Gamma]$.

The rings that arise here are $R_\infty = \mathbb{Z}[\mathbb{Q}_{>0}^\times]$, corresponding to actions of $\mathbb{Q}_{>0}^\times$, and $R_{\mathbb{Q}^d} = \mathbb{Z}[\mathbb{Q}^d]$, corresponding to actions of $\mathbb{Q}^d$. Notice that these are wildly different rings: for example, R_∞ has infinite Krull dimension, while $R_{\mathbb{Q}^d}$ has Krull dimension $d+1$. Both are non-Noetherian rings.

2. ACTIONS OF $\mathbb{Q}^d$

The main result of [12] says that for an algebraic $\mathbb{Z}^d$ -action α on a *connected* group,

$$\mathcal{M}(\alpha) > 1 \implies \mathcal{M} = \infty.$$

The same property turns out to also hold for the simplest actions of $\mathbb{Q}^d$. This is shown in Theorem 2.1 below, which is stated in a slightly more general setting. The *rational rank* of an abelian group is the maximal number of elements which are linearly independent over $\mathbb{Z}$. Thus $\mathbb{Q}^d$ and $\mathbb{Z}^d$ have rational rank d , while $\mathbb{Q}^\times$ does not have finite rational rank. If Γ has rational rank d , then $R_\Gamma = \mathbb{Z}[\Gamma]$ has Krull dimension $d+1$, and may or may not be Noetherian depending on the divisibility properties of Γ .

Theorem 2.1. *Let α be an algebraic action of a countable torsion-free group Γ of finite rational rank corresponding to a cyclic module R_Γ/I with $I \cap \mathbb{Z} = \{0\}$ and I a radical ideal. Then*

$$\mathcal{M}(\alpha) > 1 \implies \mathcal{M} = \infty.$$

Proof. The result in [12] depended on a bound for the number of solutions to S -unit equations over number fields [10] with certain uniformity properties; as shown in [11, Sect.VIII.27] it is enough to have a qualitative bound for any characteristic zero field. Here we use instead the following deep result from [4].

Theorem 2.2. [EVERTSE, SCHLICKWEI & SCHMIDT] *Let $\mathbb{K}$ be an algebraically closed field of characteristic zero, and let Γ be a finitely generated multiplicative subgroup of $(\mathbb{K}^\times)^n$ with rank r . For fixed*

$$a_1, \dots, a_n \in \mathbb{K}^\times,$$

the number of solutions $(x_1, \dots, x_n) \in \Gamma$ of the equation

$$a_1 x_1 + \dots + a_n x_n = 1$$

for which no proper subsum vanishes is bounded above by

$$\exp((6n)^{3n}(r+1)).$$

Returning to the proof of Theorem 2.1, write $\alpha = \alpha_{R_\Gamma/I}$, $X = X_{R_\Gamma/I}$ and $\lambda = \lambda_X$. Assume that the action is not mixing on r sets for some $r > 1$. It follows that there are measurable sets $A_1, \dots, A_r \subset X$ and a sequence of r -tuples

$$(q_1^{(j)}, \dots, q_r^{(j)})_{j \geq 1} \in \Gamma^r$$

such that $q_s^{(j)} - q_t^{(j)} \rightarrow \infty$ as $j \rightarrow \infty$ for $s \neq t$, for which

$$\lambda\left(\alpha_{q_1^{(j)}} A_1 \cap \dots \cap \alpha_{q_r^{(j)}} A_r\right) \not\rightarrow \prod_{s=1}^r \lambda(A_s) \text{ as } j \rightarrow \infty. \quad (1)$$

By approximating the indicator functions of the sets appearing in (1) and applying the orthogonality relations for characters on X , it follows that there are non-zero elements $a_1, \dots, a_r \in R_\Gamma/I$ with the property that

$$\widehat{\alpha_{q_1^{(j)}}}(a_1) + \dots + \widehat{\alpha_{q_r^{(j)}}}(a_r) = 0 \text{ for infinitely many } j. \quad (2)$$

First assume that the ideal I is a prime ideal P . Embed R_Γ/P into a field $\mathbb{K}$ of characteristic zero (this is possible because P is prime and $P \cap \mathbb{Z} = \{0\}$); denote by

$$x \mapsto u_1^{q_{s,1}^{(j)}} \dots u_d^{q_{s,d}^{(j)}} x = \mathbf{u}^{\mathbf{q}_s^{(j)}} x$$

the automorphism of $\mathbb{K}$ defined by the automorphism $\widehat{\alpha_{\mathbf{q}_s^{(j)}}}$ of R_Γ/P (writing $\mathbf{u}^{\mathbf{q}}$ for $u_1^{q_1} \dots u_d^{q_d}$, where $\mathbf{q} = (q_1, \dots, q_d) \in \Gamma^d$). Then equation (2) implies that

$$\mathbf{u}^{\mathbf{q}_1^{(j)}} a_1 + \dots + \mathbf{u}^{\mathbf{q}_r^{(j)}} a_r = 0 \text{ for infinitely many } j \quad (3)$$

holds in $\mathbb{K}$. Rearranging, this gives an equation

$$(-a_2/a_1)\mathbf{u}^{\mathbf{q}_2^{(j)}-\mathbf{q}_1^{(j)}} + \cdots + (-a_r/a_1)\mathbf{u}^{\mathbf{q}_r^{(j)}-\mathbf{q}_1^{(j)}} = 1 \text{ for infinitely many } j. \quad (4)$$

Assume that the rational rank of Γ is $d < \infty$ and let (Γ_n) denote a sequence of subgroups with the following properties:

- for each n , $\Gamma_n \cong \mathbb{Z}^d$;
- $\Gamma_1 \subset \Gamma_2 \subset \cdots$;
- $\Gamma = \bigcup_{n \geq 1} \Gamma_n$.

Let A_n denote the set of solutions to (4) with each $\mathbf{q}_s^{(j)} \in \Gamma_n$ for which no subsum vanishes (thus A_n is a subset of the set of values of j for which (4) holds, $A_1 \subset A_2 \subset \cdots$ and $\bigcup_{n \geq 1} A_n$ is the set of all j for which (4) holds). We may assume without loss of generality that the map

$$j \mapsto (\mathbf{q}_2^{(j)} - \mathbf{q}_1^{(j)}, \dots, \mathbf{q}_r^{(j)} - \mathbf{q}_1^{(j)})$$

is injective. Since Γ_n is isomorphic to $\mathbb{Z}^d$, Theorem 2.2 applies to show that

$$|A_n| \leq \exp((6r)^{3r}(d+1)). \quad (5)$$

Any finite bound in (5) would suffice to prove the main result in [12], namely Theorem 2.1 for $\mathbb{Z}^d$ -actions. Here the additional uniformity of Theorem 2.2 is needed: The bound in (5) is independent of n , so it follows that equation (4) holds without vanishing subsum for only finitely many j . Thus there exists a set $S \subsetneq \{2, \dots, r\}$ such that

$$\sum_{s \in S} (-a_s/a_1)\mathbf{u}^{\mathbf{q}_s^{(j)}-\mathbf{q}_1^{(j)}} = 0 \text{ for infinitely many } j. \quad (6)$$

The identity (6) shows that α is not mixing on $|S| < r$ sets. Thus for any $r < \infty$,

$$\mathcal{M}(\alpha) \leq r \implies \mathcal{M}(\alpha) < r,$$

so

$$\mathcal{M}(\alpha) > 1 \implies \mathcal{M}(\alpha) = \infty.$$

This proves Theorem 2.1 when $I = P$ is prime. Assume now that I is a radical ideal and that the system corresponding to the module R_Γ/I is not mixing on r sets for some $r > 1$ but is mixing. As before, this means there is a sequence of r -tuples

$$(\mathbf{q}_1^{(j)}, \dots, \mathbf{q}_r^{(j)})_{j \geq 1} \in \Gamma^r$$

such that $\mathbf{q}_s^{(j)} - \mathbf{q}_t^{(j)} \rightarrow \infty$ as $j \rightarrow \infty$ for $s \neq t$, for which the equation

$$\mathbf{u}^{\mathbf{q}_1^{(j)}} a_1 + \cdots + \mathbf{u}^{\mathbf{q}_r^{(j)}} a_r = 0 \text{ for infinitely many } j \quad (7)$$

holds in the ring R_Γ/I . Let

$$U = \ll c, a_1, \mathbf{u}^{\mathbf{q}} - 1 \mid c \in \mathbb{Z} \setminus \{0\}, \mathbf{q} \in \Gamma^d \gg$$

where $\langle\langle A \rangle\rangle$ denotes the multiplicative group generated by A . This is a multiplicative set, and we claim that $U \cap I = \emptyset$. If, for some $\mathbf{q} \in \Gamma^d$, $\mathbf{u}^{\mathbf{q}} - 1 \in I$, then $\alpha_{R_\Gamma/I}$ is not mixing, which is excluded by hypothesis. Since I is radical, it follows that $(\mathbf{u}^{\mathbf{q}} - 1)^m \notin I$ for all $m \geq 1$. If

$$(\mathbf{u}^{\mathbf{q}} - 1)^m b \in I \text{ for some } \mathbf{q} \in \Gamma^d, m \in \mathbb{Z} \text{ and } b \notin I$$

then we must have $(\mathbf{u}^{\mathbf{q}} - 1)^m b^m \in I$. Since I is radical, this implies that $(\mathbf{u}^{\mathbf{q}} - 1)b \in I$, so in R_Γ/I

$$b + I = \mathbf{u}^{j\mathbf{q}}b + I \text{ for all } j \geq 1,$$

which again contradicts the assumption that $\alpha_{R_\Gamma/I}$ is mixing. By induction, no product of the form

$$a_1^\ell (\mathbf{u}^{\mathbf{q}_1} - 1)^{m_1} \cdots (\mathbf{u}^{\mathbf{q}_k} - 1)^{m_k}$$

can be in I . Since R_Γ/I has no additive torsion, it follows that $U \cap I = \emptyset$. By finding a maximal ideal above I in the localization of R_Γ at U , we may find a prime ideal $P \supset I$ with the property that $P \cap U = \emptyset$ (see [2, Prop. 2.11]). The equation (7) drops via the map $x \rightarrow x + P = \bar{x}$ to a non-trivial equation

$$\mathbf{u}^{\mathbf{q}_1^{(j)}} \bar{a}_1 + \cdots + \mathbf{u}^{\mathbf{q}_r^{(j)}} \bar{a}_r = 0 \text{ for infinitely many } j, \quad (8)$$

in which not all the coefficients have vanished. It follows that the sequence

$$\left(\mathbf{q}_1^{(j)}, \dots, \mathbf{q}_r^{(j)} \right)_{j \geq 1} \in \Gamma^r$$

witnesses non-mixing on r sets in the prime system corresponding to R_Γ/P , and the argument above shows that this is only possible if $\mathcal{M}(\alpha_{R_\Gamma/P}) = 1$. By [11, Th. 1.6], this would require that there be a non-mixing element in Γ , which is impossible by the choice of U . $\square$

Theorem 2.1 does not hold without the assumption that the rational rank is finite – see Theorem 3.2. It also cannot hold without the assumption of connectedness. If the cyclic module R_Γ/I has additive torsion, then there is an element $a + I$ and an integer $k > 0$ with $ka \in I$. For a sufficiently large n , $a \in R_{\Gamma_n}$, and setting $J = I \cap R_{\Gamma_n}$ induces an inclusion

$$R_{\Gamma_n}/J \subset R_\Gamma/I$$

which dualizes to show that the original action restricted to Γ_n has a factor corresponding to $\mathbb{Z}[\Gamma_n]/J$ for some ideal J with $J \cap \mathbb{Z} \neq \{0\}$. Since $\Gamma_n \cong \mathbb{Z}^d$, finite non-trivial order of mixing is possible unless there are additional conditions on the ideal by [11].

One of the most striking features of $\mathbb{Z}^d$ -actions for $d > 1$ (as opposed to actions of $\mathbb{Z}$) is that simple examples may have order of mixing satisfying

$$1 < \mathcal{M} < \infty.$$

This was first pointed out by Ledrappier [7], who showed that the $\mathbb{Z}^2$ -action α corresponding to the module $\mathbb{Z}[u_1^{\pm 1}, u_2^{\pm 1}]/\langle 2, 1 + u_1 + u_2 \rangle$ has

$$\mathcal{M}(\alpha) = 2;$$

the paper [1] gives a simple related construction for any specified order of mixing. The same construction will give algebraic $\mathbb{Q}^d$ -actions for any $d > 1$ with any specified order of mixing.

Theorem 2.1 shows that $\mathbb{Q}^d$ -actions on connected groups also behave much like $\mathbb{Z}^d$ -actions. Further evidence for the essential similarity of algebraic $\mathbb{Q}$ -actions and $\mathbb{Z}$ -actions is provided by the next result.

Proposition 2.3. *Any mixing prime algebraic $\mathbb{Q}$ -action is mixing of all orders.*

Proof. Let the action correspond to the module $R_{\mathbb{Q}}/P$ for some prime ideal P . If the action is not mixing on r sets for some $r \geq 2$ then as before we find an equation

$$u^{q_1^{(j)}} a_1 + \cdots + u^{q_r^{(j)}} a_r = 0 \text{ for infinitely many } j \quad (9)$$

which holds in some field $\mathbb{K}$ (not necessarily of characteristic zero) with

$$q_s^{(j)} - q_t^{(j)} \rightarrow \infty \text{ in } \mathbb{Q} \text{ as } j \rightarrow \infty \text{ for } s \neq t.$$

If $P \cap \mathbb{Z} \neq \{0\}$ and P contains a non-trivial polynomial, then the variable u satisfies an algebraic equation over a finite field, so must be a root of unity in $\mathbb{K}$. This precludes mixing. So $P = \langle p \rangle$ for some rational prime p , and the $\mathbb{Q}$ -action is a full $\mathbb{Q}$ shift on p symbols, so is mixing of all orders.

If $P \cap \mathbb{Z} = \{0\}$ then we are in the setting of Theorem 2.1, which shows that the system is mixing of all orders. $\square$

3. ACTIONS OF $\mathbb{Q}_{>1}^{\times}$

Actions of the much larger group $\mathbb{Q}_{>1}^{\times}$ behave quite differently. A situation that arises naturally for such actions is one in which all the interesting parts of the action are carried by a subgroup of Γ in the following sense. Suppose that the acting group is a direct sum of two subgroups,

$$\Gamma = \Gamma' \oplus \Gamma'' \quad (10)$$

and write $\gamma = \gamma' + \gamma''$ for the unique splitting of an element γ under (10). Suppose also that there is a Γ' -invariant splitting

$$(X, \mathcal{B}, \mu) \cong \prod_{\Gamma''} (X', \mathcal{B}', \mu'), \quad (11)$$

and write elements of $\prod_{\Gamma''} X'$ in the form $(x_{\beta})_{\beta \in \Gamma''}$. Finally assume that under these isomorphisms the transformation α_{γ} on X is sent to the transformation

$$(x_{\beta})_{\beta \in \Gamma''} \mapsto (y_{\beta})_{\beta \in \Gamma''}$$

where

$$y_\beta = \alpha_{\gamma'}(y_{\beta+\gamma''}).$$

Write $\bar{\alpha}$ for the image of the restriction of α to Γ' , acting on X' . That is, the subgroup Γ' acts by restricting the original action, and the subgroup Γ'' acts as a full shift. This kind of structure has been studied by Kaminski [5] in connection with the completely positive entropy property for $\mathbb{Z}^2$ -actions. A *cylinder set* in $\prod_{\Gamma''}(X', \mathcal{B}', \mu')$ is a set of the form

$$\{(x_\beta)_{\beta \in \Gamma''} \mid x_{\beta_i} = a_i \text{ for } i \in F\}$$

where F is a finite subset of Γ'' .

Lemma 3.1. $\mathcal{M}(\alpha) = \mathcal{M}(\bar{\alpha})$.

Proof. A sequence of elements in Γ' goes to infinity if and only it goes to infinity in Γ , so

$$\mathcal{M}(\alpha) \leq \mathcal{M}(\bar{\alpha}).$$

Now let $(\gamma_1^{(j)}, \dots, \gamma_r^{(j)})_{j \geq 1}$ be a sequence of r -tuples in Γ that witnesses the statement $\mathcal{M}(\alpha) < r$. This means that for every $s \neq t$,

$$\gamma_s^{(j)} - \gamma_t^{(j)} \rightarrow \infty \text{ as } j \rightarrow \infty$$

and – without loss of generality – there are cylinder sets $A_1, \dots, A_r \in \mathcal{B}$ for which

$$\mu\left(\alpha_{\gamma_1^{(j)}} A_1 \cap \dots \cap \alpha_{\gamma_r^{(j)}} A_r\right) \not\rightarrow \mu(A_1) \cdots \mu(A_r) \text{ as } j \rightarrow \infty.$$

If, for each $s \neq t$,

$$(\gamma_s^{(j)})' - (\gamma_t^{(j)})' \rightarrow \infty \text{ in } \Gamma' \text{ as } j \rightarrow \infty$$

then the sequence $((\gamma_1^{(j)})', \dots, (\gamma_r^{(j)})')_{j \geq 1}$ witnesses that $\mathcal{M}(\bar{\alpha}) < r$, so

$$\mathcal{M}(\bar{\alpha}) \leq \mathcal{M}(\alpha)$$

as required. If for some s and along some subsequence J ,

$$(\gamma_s^{(j)})'' - (\gamma_t^{(j)})'' \rightarrow \infty \text{ in } \Gamma'' \text{ as } j \rightarrow \infty \text{ in } J \text{ for all } t \neq s,$$

then, since each A_i is a cylinder set,

$$\mu\left(\alpha_{\gamma_1^{(j)}} A_1 \cap \dots \cap \alpha_{\gamma_r^{(j)}} A_r\right) = \mu(A_s) \times \mu\left(\bigcap_{t \neq s} \alpha_{\gamma_t^{(j)}} A_t\right) \text{ for large } j \text{ in } J,$$

so

$$\mu\left(\bigcap_{t \neq s} \alpha_{\gamma_t^{(j)}} A_t\right) \not\rightarrow \prod_{t \neq s} \mu(A_t) \text{ as } j \rightarrow \infty \text{ in } J.$$

Thus the sequence with term s omitted is a witness to the statement that

$$\mathcal{M}(\bar{\alpha}) < r - 1,$$

completing the proof. $\square$

Theorem 3.2. *Let α be an algebraic action of $\mathbb{Q}_{>0}^\times$ corresponding to a module R_∞/P with $P \cap \mathbb{Z} = \{0\}$ and P a prime ideal. Then if P is finitely generated,*

$$\mathcal{M}(\alpha) > 1 \implies \mathcal{M} = \infty.$$

If P is not finitely generated, the order of mixing may be finite and positive.

Proof. View R_∞ as the ring $\mathbb{Z}[u_1^{\pm 1}, \dots]$ of Laurent polynomials in countably many commuting variables (this may be done via the map $2 \mapsto u_1$, $3 \mapsto u_2$, $5 \mapsto u_3$, and so on through the primes). If P is finitely generated, then it has a set of generating polynomials which only involve the variables $u_1, \dots, u_d$ for some $d \geq 1$. Then the action α splits up as in Lemma 3.1 with $\Gamma' \cong \mathbb{Z}^d$ and the action $\bar{\alpha}$ being the $\mathbb{Z}^d$ -action corresponding to the module $\mathbb{Z}[u_1^{\pm 1}, \dots, u_d^{\pm 1}]/P \cap \mathbb{Z}[u_1^{\pm 1}, \dots, u_d^{\pm 1}]$. By Lemma 3.1,

$$\mathcal{M}(\alpha) = \mathcal{M}(\bar{\alpha}),$$

and by [12], $\mathcal{M}(\bar{\alpha}) > 1 \implies \mathcal{M}(\bar{\alpha}) = \infty$.

For the second statement, consider the natural action of $\mathbb{Q}_{>0}^\times$ on $\widehat{\mathbb{Q}}$. This is mixing, since the equation $ax + b = 0$ in $\mathbb{Q}$ determines x . On the other hand, the action is not mixing on 3 sets since

$$1 \cdot (-1) + n \cdot (1) + (n-1) \cdot (1) = 0$$

for all n in $\mathbb{Q}$, and as elements of $\mathbb{Q}_{>0}^\times$, n , $n-1$ and $\frac{n}{n-1}$ all go to infinity and move apart as $n \rightarrow \infty$. $\square$

REFERENCES

- [1] M. Einsiedler and T. Ward. Asymptotic geometry of non-mixing sequences. *Ergodic Theory Dynam. Systems*, 23(1):75–85, 2003.
- [2] D. Eisenbud. *Commutative algebra*, volume 150 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1995. With a view toward algebraic geometry.
- [3] J.-H. Evertse. On sums of S -units and linear recurrences. *Compositio Math.*, 53(2):225–244, 1984.
- [4] J.-H. Evertse, H. P. Schlickewei, and W. M. Schmidt. Linear equations in variables which lie in a multiplicative group. *Ann. of Math. (2)*, 155(3):807–836, 2002.
- [5] B. Kamiński. Mixing properties of two-dimensional dynamical systems with completely positive entropy. *Bull. Acad. Polon. Sci. Sér. Sci. Math.*, 28(9-10):453–463 (1981), 1980.
- [6] B. Kitchens and K. Schmidt. Automorphisms of compact groups. *Ergodic Theory Dynam. Systems*, 9(4):691–735, 1989.
- [7] F. Ledrappier. Un champ markovien peut être d’entropie nulle et mélangeant. *C. R. Acad. Sci. Paris Sér. A-B*, 287(7):A561–A563, 1978.
- [8] S. MacLane. Duality for groups. *Bull. Amer. Math. Soc.*, 56:485–516, 1950.
- [9] D. W. Masser. Mixing and equations over groups in positive characteristic. *Israel J. Math.*, 142:189–204, 2004.
- [10] H. P. Schlickewei. S -unit equations over number fields. *Invent. Math.*, 102(1):95–107, 1990.

- [11] K. Schmidt. *Dynamical systems of algebraic origin*. Birkhäuser Verlag, Basel, 1995.
- [12] K. Schmidt and T. Ward. Mixing automorphisms of compact groups and a theorem of Schlickewei. *Invent. Math.*, 111(1):69–76, 1993.
- [13] A. J. van der Poorten and H. P. Schlickewei. Additive relations in fields. *J. Austral. Math. Soc. Ser. A*, 51(1):154–170, 1991.

E-mail address: `r.miles@uea.ac.uk`

SCHOOL OF MATHEMATICS, UNIVERSITY OF EAST ANGLIA, NORWICH NR4
7TJ, ENGLAND

E-mail address: `t.ward@uea.ac.uk`